

Chaos Decomposition and Gap Renormalization of Brownian Self-Intersection Local Times

Jinky Bornaes

Physics Department, MSU-IIT, Iligan City, The Philippines
jinky.bornaes@g.msuiit.edu.ph

Maria João Oliveira

Universidade Aberta, P 1269-001 Lisbon, Portugal
CMAF, University of Lisbon, P 1649-003 Lisbon, Portugal
oliveira@cii.fc.ul.pt, mjoliveira@ciencias.ulisboa.pt (New)

Ludwig Streit

Forschungszentrum BiBoS, Bielefeld University, D 33501 Bielefeld, Germany
CCM, University of Madeira, P 9000-390 Funchal, Portugal
streit@physik.uni-bielefeld.de

Abstract

We study the chaos decomposition of self-intersection local times and their regularization, with a particular view towards Varadhan's renormalization for the planar Edwards model.

Keywords: Edwards model, self-intersection local time, Varadhan renormalization, white noise analysis

Mathematics Subject Classifications (2010): 28C20, 41A25, 60H40, 60J55, 60J65, 82D60

1 Introduction

The self-intersection local time of d -dimensional Brownian motion, informally, is given as

$$L = \int_0^T dt_2 \int_0^{t_2} dt_1 \delta(\mathbf{B}(t_2) - \mathbf{B}(t_1)). \quad (1)$$

We shall see that, while "reasonably well defined" for $d = 1$, these local times become more and more singular as the dimension d increases. Intersections have thus been the object of extensive study by authors such as Dvoretzky, Erdős, Kakutani [6], [7], [8], Varadhan [35], Westwater [30], [31], [32], Le Gall [20], [21], Rosen [24], [25], [26], Dynkin [9], [10], [11], Watanabe [29], Yor [33], [34], Imkeller et al. [18], Albeverio et al. [1], [2]. For fractional Brownian motion there are papers e.g. by Rosen [27], Hu & Nualart [17], Grothaus et al. [15].

Apart from its intrinsic mathematical interest the self-intersection local time has played a role in constructive quantum field theory, and is a standard model in polymer physics for the self-repulsion ("excluded volume effect") of chain polymers in solvents [28].

Replacement of the Dirac delta function in (1) by a Gaussian

$$\delta_\varepsilon(x) := \frac{1}{(2\pi\varepsilon)^{d/2}} e^{-\frac{|x|^2}{2\varepsilon}}, \quad \varepsilon > 0,$$

leads to regularized local times

$$L_\varepsilon := \int_0^T dt_2 \int_0^{t_2} dt_1 \delta_\varepsilon(\mathbf{B}(t_2) - \mathbf{B}(t_1))$$

and for $d = 1$ one can show L^2 convergence w.r.t. white noise or Wiener measure space. But already for $d = 2$ this fails since the expectation of L_ε will diverge in the limit, asymptotically

$$\mathbb{E}(L_\varepsilon) \approx -\frac{T}{2\pi} \ln \varepsilon.$$

In this case it is sufficient to subtract the expectation, i.e. the centered regularized local time does have a well-defined L^2 limit:

$$L_{\varepsilon,c} := L_\varepsilon - \mathbb{E}(L_\varepsilon) \rightarrow L_c.$$

Apart from the Gaussian regularization above, others have been considered to remove the singularity at $t_1 = t_2$ in the integral (1). The "staircase regularization" avoids the line $t_1 = t_2$ as in see e.g. Bolthausen [5] (Fig. 1).




Fig. 1: Domain of integration for the staircase-regularized local time.

The widely used "gap regularization" does the same by omitting the strip $t_2 - t_1 < \Lambda$ in the integral. In the modelling of chain polymers the gap size Λ will be a "microscopic" quantity, i.e. of the order of the inter-monomer distance, more precisely the "Kuhn" or "persistence" length. It plays an important role in renormalization group calculations [28]: critical parameters are obtained from the postulate that macroscopic quantities do not depend on microscopic length scales.

2 Tools from White Noise Analysis [16]

Based on a d -tuple of independent Gaussian white noises $\boldsymbol{\omega} = (\omega_1, \dots, \omega_d)$ one defines a d -dimensional Brownian motion \mathbf{B} through

$$\mathbf{B}(t) \equiv \langle \boldsymbol{\omega}, \mathbb{1}_{[0,t]} \rangle = \int_0^t ds \boldsymbol{\omega}(s).$$

We shall use a multi-index notation

$$\mathbf{n} = (n_1, \dots, n_d), \quad n = \sum_{i=1}^d n_i, \quad \mathbf{n}! = \prod_{i=1}^d n_i!$$

and for d -tuples of Schwartz test functions $\mathbf{f} = (f_1, \dots, f_d) \in S(\mathbb{R}, \mathbb{R}^d)$,

$$\langle \mathbf{f}, \mathbf{f} \rangle = \sum_{i=1}^d \int dt f_i^2(t)$$

$$\langle F_{\mathbf{n}}, \mathbf{f}^{\otimes \mathbf{n}} \rangle = \int d^n t F_{\mathbf{n}}(t_1, \dots, t_n) \bigotimes_{i=1}^d f_i^{\otimes n_i}(t_1, \dots, t_n)$$

and similarly for $\langle : \boldsymbol{\omega}^{\otimes \mathbf{n}} :, F_{\mathbf{n}} \rangle$ where for d -tuples of white noise the Wick product $: \cdot : [16]$ generalizes to

$$: \boldsymbol{\omega}^{\otimes \mathbf{n}} := \bigotimes_{i=1}^d : \omega_i^{\otimes n_i} :$$

The vector valued white noise ω has the characteristic function

$$C(\mathbf{f}) := \mathbb{E}(e^{i\langle \boldsymbol{\omega}, \mathbf{f} \rangle}) = \int_{S^*(\mathbb{R}, \mathbb{R}^d)} d\mu(\boldsymbol{\omega}) e^{i\langle \boldsymbol{\omega}, \mathbf{f} \rangle} = e^{-\frac{1}{2}\langle \mathbf{f}, \mathbf{f} \rangle},$$

where $\langle \boldsymbol{\omega}, \mathbf{f} \rangle = \sum_{i=1}^d \langle \omega_i, f_i \rangle$ and $f_i \in S(\mathbb{R}, \mathbb{R})$.

Writing

$$(L^2) := L^2(S^*(\mathbb{R}, \mathbb{R}^d), d\mu)$$

there is the Itô-Segal-Wiener isomorphism with the Fock space of symmetric square integrable functions:

$$(L^2) \simeq \left(\bigoplus_{k=0}^{\infty} \text{Sym } L^2(\mathbb{R}^k, k! d^k t) \right)^{\otimes d}.$$

This implies the chaos expansion

$$\varphi(\boldsymbol{\omega}) = \sum_{\mathbf{n} \in \mathbb{N}_0^d} \langle : \boldsymbol{\omega}^{\otimes \mathbf{n}} :, F_{\mathbf{n}} \rangle \text{ for } \varphi \in (L^2)$$

with kernel functions $F_{\mathbf{n}}$ in Fock space.

Generalized functionals are constructed via a Gel'fand triple

$$(S) \subset (L^2) \subset (S)^*.$$

The generalized functionals in $(S)^*$ are conveniently characterized by their action on exponentials. In particular we use the

$$: \exp(\langle \boldsymbol{\omega}, \mathbf{f} \rangle) := C(\mathbf{f}) \exp(\langle \boldsymbol{\omega}, \mathbf{f} \rangle) \in (S)$$

to make the

Definition 1 *The transformation defined for all test functions $\mathbf{f} \in S(\mathbb{R}, \mathbb{R}^d)$ via the bilinear dual product on $(S)^* \times (S)$ by*

$$(S\Phi)(\mathbf{f}) = \langle\langle \Phi, : \exp(\langle \cdot, \mathbf{f} \rangle) : \rangle\rangle$$

is called the S -transform of $\Phi \in (S)^$.*

The multilinear expansion of $S(\Phi)$

$$(S\Phi)(\mathbf{f}) = \sum_{\mathbf{n} \in \mathbb{N}_0^d} \langle \boldsymbol{\varphi}_{\mathbf{n}}, \mathbf{f}^{\otimes \mathbf{n}} \rangle$$

extends the chaos expansion to $\Phi \in (S)^*$, with distribution valued kernels $\boldsymbol{\varphi}_{\mathbf{n}}$, such that

$$\langle\langle \Phi, F \rangle\rangle = \sum_{\mathbf{n} \in \mathbb{N}_0^d} \mathbf{n}! \langle \boldsymbol{\varphi}_{\mathbf{n}}, F_{\mathbf{n}} \rangle.$$

Definition 2 *We shall indicate the projection onto chaos of order $n \geq k$ by a superscript (k) :*

$$\langle\langle \Phi^{(k)}, F \rangle\rangle = \sum_{\mathbf{n}: n \geq k} \mathbf{n}! \langle \boldsymbol{\varphi}_{\mathbf{n}}, F_{\mathbf{n}} \rangle.$$

Proposition 3 [14]

$$\delta^{(2N)}(\mathbf{B}(t_2) - \mathbf{B}(t_1)) \in (S)^*$$

with even kernel functions

$$\psi_{2\mathbf{n}}(u_1, \dots, u_{2n}; t_1, t_2) = \frac{1}{\mathbf{n}!} (2\pi)^{-d/2} \left(\frac{1}{|t_2 - t_1|} \right)^{\frac{d}{2} + n} \left(-\frac{1}{2} \right)^n \prod_{k=1}^{2n} \mathbb{1}_{[t_1, t_2]}(u_k).$$

All the kernel functions with odd indices vanish.

Setting

$$\begin{aligned} v &:= \max(u_1, \dots, u_{2n}) \\ u &:= \min(u_1, \dots, u_{2n}) \end{aligned}$$

one computes [14] the kernel functions of the truncated local time $L^{(2N)}$ for $2N > d - 2$ by integration over $0 < t_1 < t_2 < T$:

$$\begin{aligned} \varphi_{2\mathbf{n}}(u_1, \dots, u_{2n}) &= \frac{(2\pi)^{-d/2}}{\mathbf{n}!} \left(-\frac{1}{2} \right)^n \int_0^T dt_2 \int_0^{t_2} dt_1 (t_2 - t_1)^{-n-d/2} \mathbb{1}_{[t_1, t_2]}^{\otimes 2n}(u_1, \dots, u_{2n}) \\ &= (-1)^n \left(\left(n + \frac{d}{2} - 1 \right) \left(n + \frac{d}{2} - 2 \right) (2\pi)^{d/2} 2^n \mathbf{n}! \right)^{-1} \cdot \Theta(u) \Theta(T - v) \cdot \\ &\quad \cdot (T^{-n-\frac{d}{2}+2} - v^{-n-\frac{d}{2}+2} - (T - u)^{-n-\frac{d}{2}+2} + (v - u)^{-n-\frac{d}{2}+2}) \end{aligned}$$

except for $2n = d = 2$ where

$$\varphi_2(u_1, u_2) = -\frac{1}{4\pi} (\ln v + \ln(T - u) - \ln(v - u) - \ln T) \cdot \Theta(u) \Theta(T - v).$$

The Heaviside function Θ here is the indicator function of the positive half line.

3 Regularizations

Replacement of the Dirac delta function by a Gaussian

$$\delta_\varepsilon(x) = \frac{1}{(2\pi\varepsilon)^{d/2}} e^{-\frac{|x|^2}{2\varepsilon}}, \quad \varepsilon > 0,$$

leads to regularized local times

$$L_\varepsilon = \int_0^T dt_2 \int_0^{t_2} dt_1 \delta_\varepsilon(\mathbf{B}(t_2) - \mathbf{B}(t_1)) \quad (2)$$

with kernel functions [14]

$$\begin{aligned} & \varphi_{\varepsilon, 2\mathbf{n}}(u_1, \dots, u_{2n}) \\ = & \frac{(2\pi)^{-d/2}}{\mathbf{n}!} \left(-\frac{1}{2}\right)^n \int_0^T dt_2 \int_0^{t_2} dt_1 (\varepsilon + |t_2 - t_1|)^{-n-d/2} \mathbb{I}_{[t_1, t_2]}^{\otimes 2n}(u_1, \dots, u_{2n}) \\ = & (-1)^n \left(\left(n + \frac{d}{2} - 1\right) \left(n + \frac{d}{2} - 2\right) (2\pi)^{d/2} 2^n \mathbf{n}! \right)^{-1} \cdot \Theta(u) \Theta(T - v) \cdot \\ & \cdot ((T + \varepsilon)^{-n-\frac{d}{2}+2} - (v + \varepsilon)^{-n-\frac{d}{2}+2} - (T - u + \varepsilon)^{-n-\frac{d}{2}+2} + (v - u + \varepsilon)^{-n-\frac{d}{2}+2}), \\ & \varphi_{\varepsilon, 2}(u_1, u_2) \\ = & -\frac{1}{4\pi} (\ln(v + \varepsilon) + \ln(T - u + \varepsilon) - \ln(v - u + \varepsilon) - \ln(T + \varepsilon)) \cdot \Theta(u) \Theta(T - v). \end{aligned}$$

3.1 Gap Regularization of Kernel Functions

In renormalization group studies of self-repelling Brownian motion another regularization is often used, see e.g. [28] and the references there; it suppresses intersection in small time intervals $t_2 - t_1$ between intersections by setting informally

$$L(\Lambda) := \int_{\substack{0 < t_1 < t_2 < T \\ t_2 - t_1 > \Lambda}} d^2t \delta(\mathbf{B}(t_2) - \mathbf{B}(t_1)).$$

The expectation of $L(\Lambda)$ is equal to

$$\mathbb{E}(L(\Lambda)) = \int_\Lambda^T dt_2 \int_0^{t_2-\Lambda} dt_1 \psi_0(t_1, t_2) = (2\pi)^{-d/2} \int_\Lambda^T dt_2 \int_0^{t_2-\Lambda} dt_1 (t_2 - t_1)^{-d/2}.$$

We note that for in particular $d = 2$

$$\mathbb{E}(L(\Lambda)) = -\frac{T}{2\pi} \ln \Lambda + O(1). \quad (3)$$

Recall [14] that for $\Lambda = 0$ the kernel functions $\varphi_{2\mathbf{n}}$ of the truncated local time $L^{(2N)}$ are obtained by integrating the kernel functions $\psi_{2\mathbf{n}}(u_1, \dots, u_{2n}; t_1, t_2)$

over the rectangle $0 < t_1 < u$ and $v < t_2 < T$, shaded grey in see Fig. 2. For $\Lambda > 0$ the integration is further restricted to the light domain with $t_2 - t_1 > \Lambda$. This restriction is non-trivial when $v - u < \Lambda$.

$$L^{(2N)} - L^{(2N)}(\Lambda)$$

thus has kernel functions $\rho_{2\mathbf{n}}(u, v)$ with support on

$$\{0 < u < v < T\} \cap \{v - u < \Lambda\}.$$

Fig. 2 is pertinent to the case where $\Lambda < v$ and $u < T - \Lambda$. In this case the kernel functions $\rho_{2\mathbf{n}}(u, v)$ are obtained by integrating the $\psi_{2\mathbf{n}}(u_1, \dots, u_{2n}; t_1, t_2)$ with respect to the t_i over $v - \Lambda < t_1 < u$ and $v < t_2 < t_1 + \Lambda$. Excepting the kernel function $\rho_2(u, v)$ for $d = 2$, one finds,

$$\begin{aligned} \rho_{2\mathbf{n}}(u, v) &= \int_{v-\Lambda}^u dt_1 \int_v^{t_1+\Lambda} dt_2 \psi_{2\mathbf{n}}(u_1, \dots, u_{2n}; t_1, t_2) \\ &= \frac{(2\pi)^{-d/2}}{\mathbf{n}!} \left(-\frac{1}{2}\right)^n \frac{1}{d/2 + n - 1} \cdot \\ &\quad \cdot \left(\frac{1}{d/2 + n - 2} ((v - u)^{-d/2-n+2} - \Lambda^{-d/2-n+2}) + (v - u - \Lambda)\Lambda^{-d/2-n+1} \right) \end{aligned} \quad (4)$$

and for $2n = d = 2$

$$\rho_2(u, v) = \frac{1}{4\pi} \left(\ln(v - u) - \ln \Lambda + \frac{\Lambda - v + u}{\Lambda} \right). \quad (5)$$




Fig. 2: Domain of integration for kernels of the local time, light grey for the regularized local time, dark grey for the subtraction ρ as in (4).

Using $\tau = t_2 - t_1$ we obtain the following estimate

$$\begin{aligned}
|\rho_{2\mathbf{n}}(u, v)| &= \left| \int_{v-\Lambda}^u dt_1 \int_v^{t_1+\Lambda} dt_2 \psi_{2\mathbf{n}}(u_1, \dots, u_{2n}; t_1, t_2) \right| \\
&= \frac{1}{2^n \mathbf{n}!} (2\pi)^{-d/2} \int_{v-\Lambda}^u dt_1 \int_{v-t_1}^{\Lambda} d\tau \left(\frac{1}{\tau} \right)^{\frac{d}{2}+n} \\
&\leq \frac{1}{2^n \mathbf{n}!} (2\pi)^{-d/2} \frac{1}{d/2 + n - 1} \int_{v-\Lambda}^u dt_1 (v - t_1)^{-d/2-n+1} \\
&\leq \frac{1}{2^n \mathbf{n}!} (2\pi)^{-d/2} \frac{1}{d/2 + n - 1} \frac{1}{d/2 + n - 2} (v - u)^{-d/2-n+2}
\end{aligned} \tag{6}$$

while for $d = 2n = 2$ one readily finds from (5)

$$|\rho_2(u, v)| \leq \frac{1}{4\pi} |\ln(v - u)| \tag{7}$$

when $v - u < \Lambda \ll 1$.

For very small or very large u, v , i.e. $0 < u < v < \Lambda$ or $T - \Lambda < u < v < T$, respectively, the range of integrations in (4) and (6) for $\psi_{2\mathbf{n}}(u_1, \dots, u_{2n}; t_1, t_2)$ will be $0 < t_1 < u$ and $v < t_2 < t_1 + \Lambda$, or $v < t_2 < T$ and $t_2 - \Lambda < t_1 < u$ respectively, see Fig. 3.




Fig. 3: Modified integration domains for ρ when u, v are close to zero or to T respectively.

Computations as in (4) are again straightforward, we note here only that the estimate of (6) is true also in these two cases.

Theorem 4 For $N > 0$ and $0 < \Lambda \ll T$ the chaos expansion of the gap-regularized local time $L^{(2N)}(\Lambda)$ has the kernel functions

$$\varphi_{\Lambda, 2\mathbf{n}}(u_1, \dots, u_{2n}) = \varphi_{2\mathbf{n}}(u_1, \dots, u_{2n}) - \Theta(\Lambda - (v - u))\rho_{2\mathbf{n}}(u_1, \dots, u_{2n}), \quad (8)$$

and zero otherwise, while

$$L^{(2N)} - L^{(2N)}(\Lambda)$$

has the kernel functions $\rho_{2\mathbf{n}}$ for $0 < u < v < T$, $n \geq N$.

The Heaviside function Θ exhibits the support property of the $\rho_{2\mathbf{n}}$, i.e. in the gap regularization the kernel functions of the local time are only modified when all arguments u_k are close to each other.

With these results one can in particular estimate the rate of convergence for the centered self-intersection local time in $d = 2$. Apart from the term $n = 1$ the sum

$$\|L^{(2)} - L^{(2)}(\Lambda)\|_{(L^2)}^2 = \sum_{\mathbf{n}: n \geq 1} (2\mathbf{n})! \|\rho_{2\mathbf{n}}\|_{L^2([0, T]^{2n})}^2$$

can be estimated as follows:

$$\begin{aligned} \sum_{\mathbf{n}: n > 1} (2\mathbf{n})! \|\rho_{2\mathbf{n}}\|_{L^2([0, T]^{2n})}^2 &\leq (2\pi)^{-d} \sum_{\mathbf{n}: n > 1} \frac{(2\mathbf{n})!}{(\mathbf{n}!)^2} \left(\frac{1}{2}\right)^{2n} \left(\frac{1}{d/2 + n - 1} \frac{1}{d/2 + n - 2}\right)^2 \\ &\quad \int_0^T d^{2n} u_k \left(\frac{1}{v - u}\right)^{d+2n-4} \quad v - u < \Lambda. \end{aligned}$$

We can integrate out the $2n - 2$ variables u_k with $u < u_k < v$ that lie between the smallest and the largest and obtain in this way

$$\begin{aligned} \sum_{\mathbf{n}: n > 1} (2\mathbf{n})! \|\rho_{2\mathbf{n}}\|_{L^2([0, T]^{2n})}^2 &\leq (2\pi)^{-2} \sum_{\mathbf{n}: n > 1} \frac{(2\mathbf{n})!}{(\mathbf{n}!)^2} \left(\frac{1}{2}\right)^{2n} \left(\frac{1}{n} \frac{1}{n - 1}\right)^2 \\ &\quad \cdot 2n(2n - 1) \int_0^T dv \int_0^v du \quad v - u < \Lambda \\ &\leq \frac{\Lambda T}{2\pi^2} \sum_{\mathbf{n}: n > 1} \frac{(2\mathbf{n})!}{2^{2n} (\mathbf{n}!)^2} \frac{1}{n} \frac{2n - 1}{(n - 1)^2}. \end{aligned}$$

The series is convergent (Stirling's formula). From (7) it is straightforward to estimate the remaining term with $n = 1$:

$$\|\rho_2\|_{L^2([0, T]^2)}^2 \leq \left(\frac{1}{4\pi}\right)^2 \int_0^T dt \int_0^\Lambda d\tau \ln^2 \tau = 0(T\Lambda \ln^2 \Lambda).$$

So we have shown

Theorem 5 *For $d = 2$*

$$\|L^{(2)} - L^{(2)}(\Lambda)\|_{(L^2)}^2 = 0(T\Lambda \ln^2 \Lambda) \text{ as } \Lambda \searrow 0.$$

A similar improvement of the rate of convergence has been found in the Gaussian regularization in [3].

4 Varadhan Renormalization

The model proposed by Edwards [12] for self-repelling Brownian motion suppresses self-crossings, modifying the Brownian path (or white noise) measure by a density function, informally

$$\varrho = Z^{-1} \exp(-gL)$$

with $g > 0$

$$Z = \mathbb{E}(\exp(-gL)).$$

There is no problem for $d = 1$ since L is a positive random variable and hence $\exp(-gL) < 1$. For $d = 2$ however we should replace L by the centered L_c and this then is no more positive, so that $\exp(-gL_c)$ is unbounded. The point of Varadhan's theorem is to show that this happens only on small sets so that

Theorem 6 (*Varadhan [35]*) *For $d = 2$*

$$\varrho = Z^{-1} \exp(-gL_c)$$

with

$$Z = \mathbb{E}(\exp(-gL_c))$$

is integrable.

Varadhan defines the centered local time as the limit of Gaussian approximations as in (2) and uses the Chebyshev inequality to show that $\exp(-gL_c)$ is integrable for $0 < g < \frac{\pi}{T}$.

A similar slightly stronger result can be obtained using Varadhan's technique with the approximation

$$L^{(2)}(\Lambda) \rightarrow L^{(2)} = L_c.$$

Fix $0 < \Lambda < 1$. By (3) there exists a positive constant k such that

$$L^{(2)}(\Lambda) \geq -\mathbb{E}(L(\Lambda)) \geq -k - \frac{T}{2\pi} |\ln(\Lambda)|.$$

For any constant $N \geq k + \frac{T}{2\pi} |\ln(\Lambda)|$ one has

$$\begin{aligned} \mathbb{P}(L_c \leq -N) &= \mathbb{P}(L_c - L^{(2)}(\Lambda) \leq -N - L^{(2)}(\Lambda)) \\ &\leq \mathbb{P}\left(|L^{(2)}(\Lambda) - L_c| \geq N - k - \frac{T}{2\pi} |\ln(\Lambda)|\right). \end{aligned}$$

An application of Chebyshev's inequality, using Theorem 5 then yields

$$\mathbb{P}(L_c \leq -N) \leq \frac{\mathbb{E}(|L^{(2)}(\Lambda) - L_c|^2)}{(N - k - \frac{T}{2\pi}|\ln(\Lambda)|)^2} \leq K \frac{\Lambda \ln^2 \Lambda}{(N - k - \frac{T}{2\pi}|\ln(\Lambda)|)^2}.$$

In particular, for

$$\Lambda = \exp(-\alpha(N - k)), \quad 0 < \alpha < \frac{2\pi}{T}$$

one obtains

$$\mathbb{P}(L_c \leq -N) \leq \frac{K\alpha^2}{(1 - \frac{T}{2\pi}\alpha)^2} \exp(-\alpha(N - k)).$$

Hence, $\exp(-gL_c)$ is integrable for $g < \frac{2\pi}{T}$.

For the Gaussian regularization an analogous result can be found in [3].

5 Concluding Remarks

[13], [14], [17], [18], [19], [22], [23], in the present context of regularizations and the rate of convergence see in particular [3]. Yet another regularization of the self-intersection local time is suggested by (8), namely

$$L_\Lambda^{(2N)} := \sum_{\mathbf{n}: n \geq N} \langle : \omega^{\otimes 2\mathbf{n}} :, \varphi_{2\mathbf{n}}^{(\Lambda)} \rangle$$

with kernel functions

$$\varphi_{2\mathbf{n}}^{(\Lambda)}(u_1, \dots, u_{2n}) := \Theta(v - u - \Lambda) \varphi_{2\mathbf{n}}(u_1, \dots, u_{2n})$$

where simply the range $v - u < \Lambda$ is cut out. Details of this regularization will be discussed elsewhere.

Acknowledgments

This work was financed by Portuguese national funds through FCT - Fundação para a Ciência e Tecnologia, within the project PTDC/MAT-STA/1284/2012.

References

- [1] S. Albeverio, M. J. Oliveira, L. Streit: Intersection Local Times of Independent Brownian Motions as Generalized White Noise Functionals, *Acta Appl. Math.* **69** (2001), 221–241.
- [2] S. Albeverio , Y. Z. Hu , M. Röckner and X. Y. Zhou: Stochastic Quantization of the Two-Dimensional Polymer Measure. *Appl. Math. Optim.* **40** (1996), 341–354.
- [3] W. Bock, M. J. Oliveira, J. L. Silva, and L. Streit: Polymer measure: Varadhan’s renormalization revisited. *Rev. Math. Phys.* **27** (3) (2015), 1550009.
- [4] E. Bolthausen: On the construction of the three-dimensional polymer measure. *Probab. Theory Rel. Fields* **97** (1993), 81–101.
- [5] E. Bolthausen: Large deviations and interacting random walks. *École d’Été St. Flour 1999, Lecture Notes in Math.* **1741**. Springer, Berlin, 2000.
- [6] A. Dvoretzky, P. Erdős and S. Kakutani: Double points of paths of Brownian motion in the plane, *Bull. Res. Council Israel Sect.* **F3** (1954), 364–371.
- [7] A. Dvoretzky, P. Erdős and S. Kakutani: Double points of paths of Brownian motion in n-space, *Acta Sci. Math. Szeged* **12** (1950), 75–81.
- [8] A. Dvoretzky, P. Erdős, S. Kakutani and S. J. Taylor: Triple points of the Brownian motion in 3-space, *Proc. Cambridge Philos. Soc.* **53** (1957), 856–862.
- [9] E. B. Dynkin: Polynomials of the occupation field and related random fields, *J. Funct. Anal.* **58** (1984), 20–52.
- [10] E. B. Dynkin: Self-intersection gauge for random walks and for Brownian motion, *Ann. Probab.* **16** (1988), 1–57.
- [11] E. B. Dynkin: Regularized self-intersection local times of planar Brownian motion, *Ann. Probab.* **16** (1988), 58–73.

- [12] S. F. Edwards: The statistical mechanics of polymers with excluded volume. *Proc. Roy. Soc.* **85** (1965), 613–624.
- [13] M. de Faria, C. Drumond and L. Streit: The Renormalization of Self Intersection Local Times I: The Chaos Expansion, *Inf. Dimens. Anal. Quantum Probab. Rel. Top.* **2** (2000), 223–236.
- [14] M. de Faria M., T. Hida, L. Streit and H. Watanabe: Intersection local times as generalized white noise functionals, *Acta Appl. Math.* **46** (1997), 351–362.
- [15] M. Grothaus, M. J. Oliveira, J. L. Silva and L. Streit: Self-avoiding fractional Brownian motion - The Edwards model. *J. Stat. Phys.* **145** (2011), 1513–1523.
- [16] T. Hida, H. H. Kuo, J. Potthoff and L. Streit: *White Noise - An Infinite Dimensional Calculus*. Kluwer-Academic, 1993.
- [17] Y. Hu and D. Nualart: Renormalized self-intersection local time for fractional Brownian motion. *Ann. Probab.* **33** (2005), 948–983.
- [18] P. Imkeller, V. Perez-Abreu and J. Vives: Chaos expansions of double intersection local times of Brownian motion in \mathbb{R}^d and renormalization, *Stoch. Proc. Appl.* **56** (1995), 1–34.
- [19] R. Jenane, R. Hachaichi and L. Streit: Renormalisation du temps local des points triples du mouvement Brownien, *Inf. Dimens. Anal. Quantum Probab. Rel. Top.* **9** (2006), 547–566.
- [20] J. F. Le Gall: Sur le temps local d’intersection du mouvement Brownien plan et la methode de renormalisation de Varadhan, *Sem. de Prob. XIX*, 1983/84, *Lecture Notes in Math.* **1123**, pages 314–331, Springer, Berlin 1985.
- [21] J. F. Le Gall: Sur la saucisse de Wiener et les points multiples du mouvement brownien, *Ann. Probab.* **14** (1986), 1219–1244.
- [22] S. Mendonça and L. Streit: Multiple Intersection Local Times in Terms of White Noise. *Inf. Dimens. Anal. Quantum Probab. Rel. Top.* **4** (2001), 533–543.

- [23] D. Nualart and J. Vives: Chaos expansion and local times, *Publ. Math.* **36** (1992), 827–836.
- [24] J. Rosen: A local time approach to the self-intersections of Brownian paths in space, *Comm. Math. Phys.* **88** (1983), 327–338.
- [25] J. Rosen: Tanaka’s formula and renormalisation for intersections of planar Brownian motion, *Ann. Probab.* **14** (1986), 1225–1251.
- [26] J. Rosen: A renormalized local time for multiple intersections of planar Brownian motion, *Sem. de Prob.* XX, 1984/85, *Lecture Notes in Math.* **1204**, pages 515–531, Springer 1986.
- [27] J. Rosen: The intersection local time of fractional Brownian motion in the plane. *J. Multivariate Anal.* **23** (1987), 37–46.
- [28] L. Schaefer: *Excluded Volume Effects in Polymer Solutions as Explained by the Renormalization Group*. Springer, Berlin, Heidelberg, New York, 1999.
- [29] H. Watanabe: The local time of self-intersections of Brownian motions as generalized Brownian functionals, *Lett. Math. Phys.* **23** (1991), 1–9.
- [30] M. J. Westwater: Edwards’ model for long polymer chains, *Comm. Math. Phys.* **72** (1980), 131–174.
- [31] J. Westwater: On Edwards’ model for polymer chains. II. The self-consistent potential. *Comm. Math. Phys.* **79** (1) (1981), 53–73.
- [32] J. Westwater: On Edwards’ model for polymer chains. III. Borel summability. *Comm. Math. Phys.* **84** (4) (1982), 459–470.
- [33] M. Yor: Compléments aux formules de Tanaka-Rosen. *Sem. de Prob.* XIX, 1983/84, *Lecture Notes in Math.* **1123**, pages 332–348. Springer, Berlin 1985.
- [34] M. Yor: Renormalisation et convergence en loi pour les temps locaux d’intersection du mouvement Brownien dans \mathbb{R}^3 , *Sem. de Prob.* XIX 1983/84, *Lecture Notes in Math.* **1123**, pages 350–365. Springer, 1985.
- [35] S. R. S. Varadhan: Appendix to Euclidian quantum field theory by K. Szymanzik, in: R. Jost ed., *Local Quantum Theory*. Academic Press, New York 1969.